

Weak Law of Large Numbers

- Consider I.I.D. random variables X_1, X_2, \dots
 - X_i have distribution F with $E[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$
 - Let $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
 - For any $\varepsilon > 0$:

$$P(|\bar{X} - \mu| \geq \varepsilon) \xrightarrow{n \rightarrow \infty} 0$$

- Proof:

$$E[\bar{X}] = E\left[\frac{X_1 + X_2 + \dots + X_n}{n}\right] = \mu \quad \text{Var}(\bar{X}) = \text{Var}\left(\frac{X_1 + X_2 + \dots + X_n}{n}\right) = \frac{\sigma^2}{n}$$

- By Chebyshev's inequality:

$$P(|\bar{X} - \mu| \geq \varepsilon) \leq \frac{\sigma^2}{n\varepsilon^2} \xrightarrow{n \rightarrow \infty} 0$$

Strong Law of Large Numbers

- Consider I.I.D. random variables X_1, X_2, \dots
 - X_i have distribution F with $E[X_i] = \mu$
 - Let $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$

$$P\left(\lim_{n \rightarrow \infty} \left(\frac{X_1 + X_2 + \dots + X_n}{n}\right) = \mu\right) = 1$$

- Strong Law \Rightarrow Weak Law, but not vice versa
- Strong Law implies that for any $\varepsilon > 0$, there are only a finite number of values of n such that condition of Weak Law: $|\bar{X} - \mu| \geq \varepsilon$ holds.

Intuitions and Misconceptions of LLN

- Say we have repeated trials of an experiment
 - Let event E = some outcome of experiment
 - Let $X_i = 1$ if E occurs on trial i , 0 otherwise
 - Strong Law of Large Numbers (Strong LLN) yields:

$$\frac{X_1 + X_2 + \dots + X_n}{n} \rightarrow E[X] = P(E)$$

- Recall first week of class: $P(E) = \lim_{n \rightarrow \infty} \frac{n(E)}{n}$
- Strong LLN justifies "frequency" notion of probability
- Misconception arising from LLN:
 - Gambler's fallacy: "I'm due for a win"
 - Consider being "due for a win" with repeated coin flips...

La Loi des Grands Nombres

- History of the Law of Large Numbers
 - 1713: Weak LLN described by Jacob Bernoulli



- 1835: Poisson calls it "La Loi des Grands Nombres"
 - That would be "Law of Large Numbers" in French
- 1909: Émile Borel develops Strong LLN for Bernoulli random variables
- 1928: Andrei Nikolaevich Kolmogorov proves Strong LLN in general case
- 2011: Another year passes in which Charlie Sheen does not make use of LLN
 - I'm still holding out hope for 2012...



Silence!!



And now a moment of silence...

...before we present...

...the greatest result of probability theory!

The Central Limit Theorem (CLT)

- Consider I.I.D. random variables X_1, X_2, \dots
 - X_i have distribution F with $E[X_i] = \mu$ and $\text{Var}(X_i) = \sigma^2$

$$\frac{X_1 + X_2 + \dots + X_n - n\mu}{\sigma\sqrt{n}} \rightarrow N(0, 1) \quad \text{as } n \rightarrow \infty$$

- More intuitively:

Demo

- Let $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$

- Central Limit Theorem: $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ as $n \rightarrow \infty$

- Now let $Z = \frac{\bar{X} - \mu}{\sqrt{\sigma^2/n}}$, noting that $Z \sim N(0, 1)$:

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right) \Leftrightarrow Z = \frac{\frac{1}{n}(\sum_{i=1}^n X_i) - \mu}{\sqrt{\sigma^2/n}} = \frac{n\left[\frac{1}{n}(\sum_{i=1}^n X_i) - \mu\right]}{n\sqrt{\sigma^2/n}} = \frac{(\sum_{i=1}^n X_i) - n\mu}{\sigma\sqrt{n}}$$

No Limits for Central Limit Theorem

- History of the Central Limit Theorem

- 1733: CLT for $X \sim \text{Ber}(1/2)$ postulated by Abraham de Moivre



- 1823: Pierre-Simon Laplace extends de Moivre's work to approximating $\text{Bin}(n, p)$ with Normal

- 1901: Aleksandr Lyapunov provides precise definition and rigorous proof of CLT



- 2003: Charlie Sheen stars in television series "Two and Half Men"



- By end of the 7th season (last year), there were 161 episodes
- Mean quality of subsamples of episodes is Normally distributed (thanks to the Central Limit Theorem)

Central Limit Theorem in Real World

- CLT is why many things in "real world" appear Normally distributed

- Many quantities are sum of independent variables
- Exams scores
 - Sum of individual problems
- Election polling
 - Ask 100 people if they will vote for candidate X ($p_1 = \# \text{"yes"}/100$)
 - Repeat this process with different groups to get p_1, \dots, p_n
 - Have a normal distribution over p_i
 - Can produce a "confidence interval"
 - How likely is it that estimate for true p is correct
 - We'll do an example like that soon

This is Your Midterm on the CLT

- Start with 180 midterm scores: X_1, X_2, \dots, X_{180}

- $E[X_i] = 68.9$ and $\text{Var}(X_i) = 611.37$


- Created 18 disjoint samples of size $n = 10$

- $Y_1 = \{X_1, \dots, X_{10}\}, Y_2 = \{X_{11}, \dots, X_{20}\}, Y_i = \{X_{10(i-1)+1}, \dots, X_{10i}\}$

$$\bar{Y}_i = \frac{1}{10} \sum_{j=1}^{10} Y_{ij}$$

- Prediction by CLT: $\bar{Y}_i \sim N(68.9, 611.37/18 \approx 33.965)$

$$Z_i = \frac{\bar{Y}_i - E[X_i]}{\sqrt{\sigma^2/n}} = \frac{\bar{Y}_i - 68.9}{\sqrt{611.37/18}} \quad \bar{Z} = \frac{1}{18} \sum_{i=1}^{18} Z_i = 8.2 \times 10^{-16} \quad \text{Var}(\bar{Z}) = 0.914$$



Estimating Clock Running Time

- Have new algorithm to test for running time

- Mean (clock) running time: $\mu = t$ sec.
- Variance of running time: $\sigma^2 = 4 \text{ sec}^2$.
- Run algorithm repeatedly (i.i.d. trials), measure time
 - How many trials so estimated time = $t \pm 0.5$ with 95% certainty?
 - $X_i =$ running time of i -th run (for $1 \leq i \leq n$)
 - By Central Limit Theorem, $Z \sim N(0, 1)$, where:

$$Z_n = \frac{(\sum_{i=1}^n X_i) - n\mu}{\sigma\sqrt{n}} = \frac{(\sum_{i=1}^n X_i) - nt}{2\sqrt{n}}$$

$$P(-0.5 \leq \frac{\sum_{i=1}^n X_i}{n} - t \leq 0.5) = P(-\frac{0.5\sqrt{n}}{2} \leq \frac{\sum_{i=1}^n X_i - nt}{2\sqrt{n}} \leq \frac{0.5\sqrt{n}}{2}) = P(-\frac{0.5\sqrt{n}}{2} \leq Z_n \leq \frac{0.5\sqrt{n}}{2})$$

$$= \Phi(\frac{\sqrt{n}}{4}) - \Phi(-\frac{\sqrt{n}}{4}) = \Phi(\frac{\sqrt{n}}{4}) - (1 - \Phi(\frac{\sqrt{n}}{4})) = 2\Phi(\frac{\sqrt{n}}{4}) - 1 \approx 0.95 \Rightarrow \Phi(\frac{\sqrt{n}}{4}) = 0.975$$

- Solve for n^* : $\frac{\sqrt{n^*}}{4} = 1.96 \Rightarrow n^* = \lceil (7.84)^2 \rceil = 62$

Estimating Time With Chebyshev

- Have new algorithm to test for running time

- Mean (clock) running time: $\mu = t$ sec.

- Variance of running time: $\sigma^2 = 4 \text{ sec}^2$.

- Run algorithm repeatedly (i.i.d. trials), measure time

- How many trials so estimated time = $t \pm 0.5$ with 95% certainty?

- $X_i =$ running time of i -th run (for $1 \leq i \leq n$)

- What would Chebyshev say? $P(|X_S - \mu_S| \geq k) \leq \frac{\sigma_S^2}{k^2}$

$$\mu_S = E\left[\sum_{i=1}^n \frac{X_i}{n}\right] = t \quad \sigma_S^2 = \text{Var}\left(\sum_{i=1}^n \frac{X_i}{n}\right) = n \frac{\sigma^2}{n^2} = \frac{4}{n}$$

$$P\left(\left|\sum_{i=1}^n \frac{X_i}{n} - t\right| \geq 0.5\right) \leq \frac{4/n}{(0.5)^2} = \frac{16}{n} = 0.05 \Rightarrow n \geq 320$$

- Thanks for playing Pafnuty...

Crashing Your Web Site

- Number visitors to web site/minute: $X \sim \text{Poi}(100)$

- Server crashes if ≥ 120 requests/minute

- What is $P(\text{crash in next minute})$?

- Exact solution: $P(X \geq 120) = \sum_{i=120}^{\infty} \frac{e^{-100}(100)^i}{i!} \approx 0.0282$

- Use CLT, where $\text{Poi}(100) \sim \sum_{i=1}^n \text{Poi}(100/n)$ (all i.i.d.)

$$P(X \geq 120) = P(X \geq 119.5) = P\left(\frac{X - 100}{\sqrt{100}} \geq \frac{119.5 - 100}{\sqrt{100}}\right) = 1 - \Phi(1.95) \approx 0.0256$$

- Note: Normal can be used to approximate Poisson

- I'll give you one more chance (one-sided) Chebyshev:

$$P(X \geq 120) = P(X \geq E[X] + a) \leq \frac{\sigma^2}{\sigma^2 + a^2} = \frac{100}{100 + 20^2} = 0.2$$



It's play time!

Sum of Dice

- You will roll 10 6-sided dice (X_1, X_2, \dots, X_{10})
 - X = total value of all 10 dice = $X_1 + X_2 + \dots + X_{10}$
 - Win if: $X \leq 25$ or $X \geq 45$
 - Roll!

• And now the truth (according to the CLT):
 $E[X] = 10E[X_i] = 10(3.5) = 35$ $\text{Var}(X) = 10 \text{Var}(X_i) = 10 \frac{35}{12} = \frac{350}{12}$

$$1 - P(25.5 \leq X \leq 44.5) = 1 - P\left(\frac{25.5 - 35}{\sqrt{350/12}} \leq \frac{X - 35}{\sqrt{350/12}} \leq \frac{44.5 - 35}{\sqrt{350/12}}\right)$$
$$\approx 1 - (2\Phi(1.76) - 1) \approx 2(1 - 0.9608) = 0.0784$$

- If only Chebyshev were right...

$$P(|X - \mu| \geq k) = P(|X - 35| \geq 10) \leq \frac{\sigma^2}{k^2} = \frac{350/12}{100} \approx 0.292$$